

Sensor technology for the identification of mycotoxins in cereals

Claudia Rasch, Dörte Steinbrück, Michael U. Kumke, Hans-Gerd Löhmannsröben

University of Potsdam, Department of Chemistry (Physical Chemistry)

Karl-Liebknecht-Str. 24-25, 14476 Potsdam-Golm (Germany)

Mycotoxins are toxic fungal metabolites that may contaminate primary food products such as cereals, nuts and fruits. The most predominant mycotoxins in Europe are among others the *Aflatoxins* and *Ochratoxins* produced by storage fungi, such as *Aspergillus* and *Penicillium* species, and Toxins from field-borne *Fusarium* species, for example *Zearalenone* and *Deoxynivalenol*.

Because of the potential health hazards for humans, the monitoring of food and feed for the presence of these toxins are highly important with respect to the food safety aspects. Therefore, an urgent need exists for reliable, low-cost and easy-to-handle methods for the detection of fungi contamination and the measurement of the mycotoxin content in-situ especially in the raw materials at the beginning of the food production chain.

The use of spectroscopic methods in the food research is increasing, in particular in combination with chemometric tools. Non-destructive methods, such as absorption, fluorescence and reflection spectroscopy, are powerful methods for the detection of mycotoxins in solution and on the surface of grains and flour. The introduction of fiber-based chemical sensors (FOCS) might further accelerate the application of spectroscopic methods in the field of feed and food control.

Because of its outstanding sensitivity fluorescence spectroscopy based techniques are especially suited for the in-situ detection of mycotoxins such as *Ochratoxins*, *Aflatoxins* as well as *Zearalenone*. These mycotoxins can be monitored with an excitation in the spectral range of $200 \text{ nm} < \lambda_{\text{ex}} < 400 \text{ nm}$ and a detection wavelength $\lambda_{\text{em}} > 420 \text{ nm}$. In combination with absorption and reflection measurements, qualitative and quantitative information on the mycotoxins present can be obtained. In addition, the NIR reflection spectra yield further information on the ingredients, the moisture content, and the presence (or absence) of fungi in the sample.